



COMPOSITION

| Levamisole (sf hydrochloride) | . 20 g |
|-------------------------------|--------|
| Excipient qs | 100 g |

PHARMACEUTICAL PROPERTIES

Levamisole is an anthelmintic from the imidazothiazole family with great activity on the larval and adult forms of gastrointestinal and pulmonary strongles. Its good digestive bioavailability and its excellent tissue diffusion allow it to reach pulmonary nematodes and forms in hypobiosis.

TARGET SPECIES

Poultry, sheep, goats, cattle.

INDICATIONS

In poultry: treatment of capillariosis, heterakidosis, ascariasis. In ruminants: treatment of gastrointestinal and pulmonary strongyloses.

ADMINISTRATION AND DOSAGE

Oral use. Poultry: In drinking water at a dose of 20 mg of active ingredient per kg of bodyweight or 10 g of powder / 100 kg of bodyweight. Ruminants: 8 mg of active ingredient / kg or 0.40 g of powder / 10 kg of live weight. Dissolve the contents of a 100 g sachet in 1 liter of water, shake well and administer 4 ml / 10 kg bodyweight.

CONTRAINDICATIONS AND INTERACTIONS

- Do not use in dairy females whose milk is intended for human consumption and in laying hens whose eggs are intended for human consumption.
- Do not use in equines.
- Do not combine with organophosphates and specialties containing diethylcarbamazine citrate.



SIDE EFFECTS

Mild and transient signs such as restlessness, urination, defecation and ptyalism are sometimes seen after treatment with levamisole preparations.

PRECAUTIONS FOR USE Keep the product out of reach of children

WITHDRAWAL PERIOD Meat and offal: 7 days.

STORAGE CONDITIONS Keep the product in the original packaging, protect from light and at a temperature $\leq 25 \degree \text{C}$.

PRESENTATIONS 100 g sachets.