

ATLAPRIM ORAL SOLUTION

COMPOSITION

Sulfadiazine (sf sodium)	g
Trimethoprim	g
Excipient qs 100 r	nl

PHARMACEUTICAL PROPERTIES

The TMP-SDZ association is known for its synergistic effects against many pathogenic germs. This synergy results in a broadening of the spectrum which includes germs insensitive to the two products taken separately, a powerful bactericidal activity, while the two products are bacteriostatic, a weak appearance of resistant germs. The two antibacterials TMP and sulfadiazine have fairly similar pharmacokinetic properties: good digestive absorption and good tissue diffusion.

TARGET SPECIES

Poultry and pre-ruminant calf

INDICATIONS

Infections with germs sensitive to the Sulfamide-Trimethoprim combination in the target species, ATLAPRIM is particularly indicated in the treatment of colibacillosis, salmonellosis, respiratory infections, infectious enteritis and staphylococcal disease

ADMINISTRATION AND DOSAGE

- Poultry: 30 mg / kg bodyweight or 30 ml solution / 100 kg bodyweight in drinking water once a day for 5 consecutive days.
- Calf: 30 mg / kg / day for 5 days, i.e. 3 ml / 10 kg bodyweight per day for 5 days.

The daily dose can be administered in one or two divided doses.

The product can be administered after dilution in water or in milk replacer.

CONTRAINDICATIONS AND INTERACTIONS

- Serious functional disorders of the liver and kidney,
- Hypersensitivity to sulfonamides.
- Administration to laying hens and dairy females whose products are intended for human consumption.



SIDE EFFECTS

Under normal conditions of use, the combination TMP-sulfadiazine is well tolerated.

PRECAUTIONS FOR USE

- Respect the indicated dosage.
- Ensure sufficient water ration. Keep out of the reach of children.

WITHDRAWAL PERIOD

Meat and offal: - 12 days

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature \leq 25 $^{\circ}$ C.

PHARMACEUTICAL FORM AND PRESENTATION

250 ml, 500 ml and 1 liter bottles.