

DIARICUR ORAL POWDER

COMPOSITION

| Sulfaguanidine | 90.00 g |
|----------------|---------|
| Excipient qs | 100 g |

PHARMACEUTICAL PROPERTIES

Sulfaguanidine, an antibacterial in the sulfonamide family, is active on most bacteria responsible for gastroenteritis. It is a bacteriostatic antibiotic. Its mode of action results in the inhibition of the synthesis of folic acid. The pharmacokinetics of sulfaguanidine is characterized by very low absorption in the digestive tract where it remains in active form, which makes it a product of choice for the treatment of infectious gastroenteritis.

TARGET SPECIES

Ruminants, Poultry, Horses.

INDICATIONS

Infectious gastroenteritis, Diarrhea of infectious origin.

ADMINISTRATION AND DOSAGE

Oral use after dilution in drinking water the dose of:

Young calves: 1 sachet / day in two divided doses. Heavy calves and foals: 2 sachets / day in two doses.

Lambs and kids: ½ sachet / subject in two catches.

Poultry: 10 sachets / 100 liters of water once a day.

Large animals: 4-5 sachets / day divided into 3 doses. The duration of treatment is 3 to 5

consecutive days.



CONTRAINDICATIONS AND INTERACTIONS

Hypersensitivity to sulfonamides. Renal failure.

Beneficial interactions with polymyxin-colistin, macrolides.

Do not combine with nephrotoxic products, bismuth, mineral oils, opiates.

SIDE EFFECTS

If the doses and contraindications are respected, no side effects are observed in the target species.

PRECAUTIONS FOR USE

Consider the dosages.

- Give animals plenty to drink.
- Keep the product out of the reach of children.
- Keep the product at a temperature not exceeding 25 ° C and protect from light.

WITHDRAWAL PERIOD

Meat and offal: 3 days Milk: 2 days Eggs: 8 days

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature \leq 25 ° C.

PHARMACEUTICAL FORM AND PRESENTATION

Bag of 10 g