

HEPATOCHOL

ORAL SOLUTION

COMPOSITION

Sorbitol	50.00 g
DL-Methionine	1.50 g
Chloride choline	13, 40 g
Excipient qs	100 ml

PHARMACEUTICAL PROPERTIES

Methionine, sorbitol and choline are most often combined for their lipotropic and hepatotropic properties. Sorbitol, is a very assimilable sugar which has a detoxifying action and promotes bile, pancreatic and gastric secretions in a lasting way. Methionine, a sulfur amino acid, has a lipotropic action on the liver. Choline unites with liver triglycerides to form phospholipids, which are their forms of transport in the body to reserve fatty tissue.

TARGET SPECIES

Poultry, ruminants, horses and carnivores.

INDICATIONS

In ruminants, horses and carnivores: Hepatoprotection, hepatic failure, hepatic jaundice and poisoning.

In poultry: hepatoprotection, hepatorenal syndrome and fatty liver.

ADMINISTRATION AND DOSAGE

- Oral route for 5 to 10 consecutive days at the following doses:
- Cattle and Equine: 25 to 50 ml / day;
- Sheep, Goats and Calves: 10 to 15 ml / day;
- Poultry: 1 to 2 L / Tonne of drinking water / day;
- Dogs: 10 to 30 Kg: 5 ml / day; 2 to 10 kg: 2.5 ml / day;
- Cats: 1 ml / day



CONTRAINDICATIONS AND INTERACTIONS

None under the recommended conditions of use.

SIDE EFFECTS

If the dose is respected, no side effects are known

PRECAUTIONS FOR USE

- Take the recommended dosage;
- Keep the product out of the reach of children
- Protect from light and at a temperature below 25 $^{\circ}$ C.

WITHDRAWAL PERIOD

NONE

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature \leq 25 $^{\circ}$ C.

PHARMACEUTICAL FORM AND PRESENTATION

500 ml, 1 liter and 5-liter bottles