



ANTIBIOVIT

ORAL POWDER

COMPOSITION

Oxytetracycline (sf.chlorhydrate).....	50,00g
Vitamin A palmitate	10 000 000,00 UI
Vitamin E acetate.....	..6 000 UI
Vitamin D3.....	1 250 000,00 UI
Vitamin K3 (menadione sodium bisulfite).....	8,00 g
Vitamin C.....	22,00 g
Vitamin B2	3,00 g
Vitamin B12.....	10,00 mg
Biotine.....	4,00 mg
Excipient q.s.p.....	100 g

PHARMACEUTICAL PROPERTIES

Antibiovit is based on a broad spectrum antibacterial, Oxytetracycline, and vitamins. Oxytetracycline is characterized by its very broad spectrum of activity which includes most pathogenic germs in domestic animals. Oxytetracycline is also characterized by good oral bioavailability, a large volume of distribution allowing very high tissue concentrations to be obtained and elimination in large part in active form by the kidney and bile. The addition of vitamins in the preparation helps support the body to better fight against diseases.

TARGET SPECIES

Poultry

INDICATIONS

Conditions with germs sensitive to Oxytetracycline.

It is mainly about the prevention and treatment of the following infections:

- Chronic respiratory disease ‘MRC’,
- Infectious coryza, • Avian cholera,
- Infectious sinusitis of the turkey,
- Bacterial complications of viral infections.



ADMINISTRATION AND DOSAGE

Oral use in drinking water at a dose of 20 mg of Oxytetracycline / kg bodyweight / day corresponding to 0.40 g of Antibiovit / kg / day for 3 to 5 consecutive days. As an indication, ANTIBIOVIT can be given at a rate of 1Kg / 250 L of drinking water. For the preparation of the drug solution, make a predilution of the quantity of powder to be administered in a little water before its incorporation in the drinking water tank.

CONTRAINDICATIONS AND INTERACTIONS

- Do not use in patients with severe hepatic or renal impairment.
- Do not administer to subjects who have shown hypersensitivity to tetracyclines.
- Do not combine with digestive dressings based on aluminum or calcium salts.

SIDE EFFECTS

If the dose is respected, no side effects or side effects are known for oral Oxytetracycline.

PRECAUTIONS FOR USE

1. Adopt the prescribed dosages
2. Keep the product out of the reach of children.
3. Ensure good hydration of animals during treatment
4. Keep the product at a temperature not exceeding 25 ° C and protect from light.

WITHDRAWAL PERIOD

Meat and offal: 7 days

STORAGE CONDITIONS

Keep the product in the original packaging, protect from light and at a temperature $\leq 25^{\circ}$ C.

PHARMACEUTICAL FORM AND PRESENTATION

Sachets of 100 g and 1 Kg